




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/04, 207/40, A61K 31/55 // (C07D 487/04, 223:00, 209:00)	A1	(11) International Publication Number: WO 98/41527 (43) International Publication Date: 24 September 1998 (24.09.98)
(21) International Application Number: PCT/JP98/01085 (22) International Filing Date: 16 March 1998 (16.03.98) (30) Priority Data: 9/79246 14 March 1997 (14.03.97) JP (71) Applicant (for all designated States except US): SUNTORY LIMITED [JP/JP]; 1-40, Dojimahama 2-chome, Kita-ku, Osaka-shi, Osaka 530-0004 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): MIZUNO, Akira [JP/JP]; 17-1-1-811, Fukakusashinmonjo-cho, Fushimi-ku, Kyoto-shi, Kyoto 612-8436 (JP). SHIBATA, Makoto [JP/JP]; 946-5-506, Tanaka-cho, Ashikaga-shi, Tochigi 326-0822 (JP). IWAMORI, Tomoe [JP/JP]; 1-1-37-402, Sojiji, Ibaraki-shi, Osaka 567-0801 (JP). INOMATA, Norio [JP/JP]; 5-8-1, Sakurai, Mino-shi, Osaka 562-0043 (JP). (74) Agent: ONO, Nobuo; Kanda SK Building 6F, 3-22, KandaSakuma-cho, Chiyoda-ku, Tokyo 101-0025 (JP).		(81) Designated States: AU, CA, CN, HU, KR, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PYRROLOAZEPINE COMPOUNDS (57) Abstract Disclosed are pyrroloazepine compounds and their salts. These pyrroloazepine compounds are represented by formula (I) wherein the dashed line indicates the presence or absence of a bond; and, when the bond is present, Z ₂ is not present and Z ₁ is H but, when the bond is absent, Z ₁ and Z ₂ are both Hs; Z ₁ is H and Z ₂ is OH; Z ₁ and Z ₂ are both SR ₅ s in which R ₅ is alkyl, aralkyl or aryl; or Z ₁ and Z ₂ are combined together to represent O, NOR ₆ in which R ₆ is H, alkyl, aralkyl or aryl, or C ₂ -C ₃ alkylenedithio; R is H, alkyl, cycloalkyl, cycloalkyl alkyl or aralkyl; and the ring P is a specific pyrrole ring. These pyrroloazepine compounds and salts are effective as preventives or therapeutics for general circulatory diseases such as hypertension, heart failure, ischemic hear diseases, cerebrovascular disturbances and peripheral circulatory disturbances. Their production processes are also disclosed.		



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

DESCRIPTION**PYRROLOAZEPINE COMPOUNDS****5 Technical Field**

 This invention relates to novel pyrroloazepine compounds. More specifically, this invention is concerned with pyrrolo[3,2-c]azepine compounds, pyrrolo-[3,4-c]azepine compounds and salts thereof, said com-
10 pounds and salts having strong α_1 -blocking action and serotonin-2 receptor antagonistic action and being useful as pharmaceuticals for use in the prevention or treatment of circulatory diseases such as hypertension, heart failure, ischemic heart diseases such as angina
15 pectoris, myocardial infarction and post-PTCA restenosis, cerebrovascular disturbances such as cerebral infarction and cerebral sequelae after sub-arachnoid hemorrhage, and peripheral circulatory disturbances such as arteriosclerosis obliterans, thrombo-
20 angiitis obliterans, Raynaud disease and Buerger disease; their preparation process; and pharmaceuticals containing them as effective ingredients.

Background Art

25 As pharmaceuticals which act on the circulatory

- 2 -

system, many products are known to date, including a variety of products developed as vasodilators.

Among such vasodilators, α_1 -blockers led by prazosin are the targets of active developments, because they have advantages such that (1) their anti-hypertensive action is strong and reliable, (2) they do not adversely affect lipometabolism or saccharo-metabolism and (3) they can be used easily for hypertensives suffering from complication. As α_1 -blockers which are currently in clinical use, bunazosin, terazosin, urapidil, doxazosin and the like can be mentioned in addition to prazosin. Further, medicines having α_1 -blocking action and anti-serotonin action in combination are expected to become still better therapeutics for hypertension, because they have possibility to reduce side effects, such as orthostatic hypotension and reflex tachycardia, induced by antihypertensive action which is based on α_1 -blocking action.

Further, a hypertensive is considered to be prone to an ischemic heart disease or peripheral circulatory disturbance, since his or her platelet aggregating ability has been generally potentiated to have higher thrombophilia. As one of those taking part in thrombosis, serotonin is known. Serotonin is a compound contained abundantly in platelets, which are a blood

- 3 -

component, and in a central nervous system, on the other hand, it acts as a neurotransmitter. In platelets, it is released upon stimulation by thromboxane A₂, ADP, collagen or the like, and synergistically acts on release of various platelet aggregation factors through activation of serotonin-2 receptors in the platelets and vascular smooth muscle cells and also on vasoconstriction by norepinephrine through α_1 receptors, thereby inducing strong platelet aggregation and vasoconstriction [P.M. Vanhoutte, "Journal of Cardiovascular Pharmacology", Vol. 17 (Suppl. 5), S6-S12 (1991)].

Serotonin is also known to potentiate proliferation of vascular smooth muscle cells [S. Araki et al., "Atherosclerosis", Vol. 83, pp.29-34(1990)]. It has been considered that, particularly when endothelial cells are injured as in arteriosclerosis or myocardial infarction, the vasoconstricting action and thrombus forming action of serotonin are exasperated, thereby reducing or even stopping blood supply to myocardial, cerebral and peripheral organs [P. Golino et al., "The New England Journal of Medicine", Vol. 324, No. 10, pp.641-648(1991), Y. Takiguchi et al., "Thrombosis and Haemostasis", Vol. 68(4), pp.460-463(1992), A.S. Weyrich et al., "American Journal of Physiology", Vol.

- 4 -

263, H349-H358(1992)]. Being attracted by such actions of serotonin or serotonin-2 receptors, various attempts are now under way to use a serotonin-2 receptor antagonist as a pharmaceutical for ischemic diseases of the heart, the brain and peripheral tissues.

From the foregoing, a medicine having α_1 -blocking action and serotonin-2 receptor antagonistic action in combination is expected to have vasodilative action, antiplatelet action and vascular smooth muscle proliferation inhibiting action, and is considered to become a medicine extremely effective for the prevention or treatment of not only hypertension but also general circulatory diseases such as heart failure, ischemic heart diseases such as angina pectoris, myocardial infarction and post-PTCA restenosis, cerebrovascular disturbances such as cerebral infarction and cerebral sequelae after subarachnoid hemorrhage, and peripheral circulatory disturbances such as arteriosclerosis obliterans, thromboangiitis obliterans, Raynaud disease and Buerger disease.

Until today, several medicines have been reported to have α_1 -blocking action and serotonin-2 receptor antagonistic action in combination. They are however still accompanied with many problems to be improved in potency, selectivity to other receptors, toxicity, side


- 5 -

effects and the like. There is hence an outstanding desire for the provision of a still better compound.

Disclosure of the Invention

5 In view of the foregoing circumstances, the present inventors have proceeded with extensive research, resulting in the finding of pyrrolo[3,2-c]azepine compounds and pyrrolo[3,4-c]azepine compounds which have strong α_1 -blocking action and serotonin-2 receptor
10 antagonistic action in combination, have low toxicity and less side effects, and are useful for the prevention and treatment of general circulatory diseases such as hypertension, heart failure, ischemic heart diseases, cerebrovascular disturbances and peripheral
15 circulatory disturbances.

The present invention has been completed based on the above described findings. A first object of the present invention is to provide a pyrroloazepine compound or a salt thereof, said pyrroloazepine compound
20 being represented by the following formula (I):



wherein


- 6 -

the ring P represented by




means a pyrrole ring represented by the following structure:


5



or




in which A represents an alkylene group, an alkenylene group or an alkynylene group, and Y represents a group



in which W represents CH, C= or a nitrogen atom; and,
 10 when W represents CH, m stands for 0 or 1, B represents an oxygen atom, a sulfur atom, a carbonyl group, a sulfinyl group, a sulfonyl group, an alkylene group, an alkenylene group, a group -C(OH)R₁- in which R₁ represents a substituted or unsubstituted aryl group, a
 15 group -CHR₂- in which R₂ represents a substituted or unsubstituted aryl group, or a substituted or unsubstituted, cyclic or acyclic acetal group; when W represents C=, m stands for 1, B represents a group

- 7 -



in which the double bond is coupled with W and R₃ represents a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group; when W represents a nitrogen atom, m stands for 0 or 1, and B represents a carbonyl group, a sulfonyl group, an alkylene group, an alkenylene group or a group -CHR₄- in which R₄ represents a substituted or unsubstituted aryl group; E₁ and E₂ each independently represents a hydrogen atom or a lower alkyl group; and D represents a substituted or unsubstituted aromatic hydrocarbon group or a substituted or unsubstituted aromatic heterocyclic group;

the dashed line indicates the presence or absence of a bond; and, when the bond indicated by the dashed line is present, Z₂ is not present and Z₁ represents a hydrogen atom but, when the bond indicated by the dashed line is absent, Z₁ and Z₂ both represent hydrogen atoms; Z₁ represents a hydrogen atom and Z₂ represents a hydroxyl group; Z₁ and Z₂ both represent groups SR₅ in which R₅ represents a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group; or Z₁ and Z₂ are combined together to represent

- 8 -

an oxygen atom, a group NOR_6 in which R_6 represents a hydrogen atom, a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group, or a group

5



in which G represents a substituted or unsubstituted ethylene group or a substituted or unsubstituted trimethylene group; and

R represents a hydrogen atom, a linear or
10 branched alkyl group, a cycloalkyl group, a cycloalkyl-alkyl group or a substituted or unsubstituted aralkyl group.

A second object of the present invention is to provide a preparation process of the pyrroloazepine
15 compound (I) or its salt.

Further, a third object of the present invention is to provide a pharmaceutical which comprises the pyrroloazepine compound (I) or its pharmacologically-acceptable salt as an effective ingredient and is
20 usable for the treatment or the like of circulatory diseases.

- 9 -

Best Modes for Carrying Out the Invention

In the pyrroloazepine compound (I) of the present invention, preferred examples of the group R can include a hydrogen atom; linear or branched alkyl groups having 1-8 carbon atoms preferably, such as methyl, ethyl, n-propyl, isopropyl and n-pentyl; cycloalkyl groups having 3-8 carbon atoms, such as cyclopropyl, cyclopentyl and cyclohexyl; cycloalkyl-alkyl groups having 4-8 carbon atoms, such as cyclopropylmethyl, cyclohexylmethyl and cyclohexylethyl; and aralkyl groups having 7-22 carbon atoms, such as diphenylmethyl, benzyl and phenethyl. One or more hydrogen atoms of each of these groups may be substituted by a like number of halogen atoms such as fluorine, chlorine and/or bromine atoms, alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl, and/or alkoxy groups having 1-4 carbon atoms preferably, such as methoxy and/or ethoxy. Particularly preferred examples of the group R can be methyl and ethyl.

Further, preferred examples of the group Z_1 and the group Z_2 in the compound (I) according to the present invention can include the following combinations: when the bond indicated by the dashed line is present, Z_2 is not present and Z_1 represents a hydrogen atom; when the bond indicated by the dashed line is absent,

- 10 -

Z_1 and Z_2 are both hydrogen atoms, Z_1 and Z_2 both
 represent the groups SR_5 , Z_1 represents a hydrogen atom
 and Z_2 represents a hydroxyl group, and Z_1 and Z_2 are
 combined together to represent an oxygen atom, the
 5 group NOR_6 or the group



wherein G has the same meaning as defined above.

Preferred examples of R_6 in the group NOR_6 can
 include a hydrogen atom; linear or branched alkyl
 10 groups having 1-4 carbon atoms preferably, such as
 methyl and ethyl; aryl groups having 6-14 carbon atoms,
 such as phenyl and naphthyl; and aralkyl groups having
 7-22 carbon atoms, such as benzyl and phenethyl. One
 or more of the hydrogen atoms of each of these groups
 15 may be substituted by a like number of halogen atoms
 such as fluorine, chlorine and/or bromine atoms, alkyl
 groups having 1-4 carbon atoms preferably, such as
 methyl and/or ethyl, and/or alkoxy groups having 1-4
 carbon atoms preferably, such as methoxy and/or ethoxy.
 20 Of these, hydrogen atom and methyl group are particu-
 larly preferred.

Further, preferred examples of G in the group

- 11 -



can include ethylene and trimethylene. One or more of the hydrogen atoms of each of these groups may be substituted by a like number of halogen atoms such as

5 fluorine, chlorine and/or bromine atoms, alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl, aryl groups having 6-14 carbon atoms, such as phenyl and naphthyl, aralkyl groups having 7-22 carbon atoms, such as benzyl and phenethyl, and/or

10 alkylidene groups having 1-4 carbon atoms preferably, such as methylenidene and/or ethylenidene.

Preferred examples of R_5 in the group SR_5 can include linear or branched alkyl groups having 1-4 carbon atoms preferably, such as methyl and ethyl; aryl groups

15 having 6-14 carbon atoms such as phenyl and naphthyl; and aralkyl groups having 7-22 carbon atoms such as benzyl and phenethyl. One or more of the hydrogen atoms of each of these groups may be substituted by a like number of halogen atoms such as fluorine, chlorine

20 and/or bromine atoms, alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl, and/or alkoxy groups having 1-4 carbon atoms preferably, such as methoxy and/or ethoxy.

- 12 -


In the pyrroloazepine compounds (I) of the present invention, the ring P represents any one of the following pyrrole rings:



5 wherein A and Y have the same meanings as defined above.

Preferred examples of the group A can include linear or branched alkylene groups having 2-10 carbon atoms, such as ethylene, trimethylene, tetramethylene, pentamethylene and octamethylene; linear or branched alkenylene groups having 4-10 carbon atoms, such as 2-butenylene and 3-pentenylene; and linear or branched alkynylene groups having 4-10 carbon atoms, such as 2-butyneylene and 3-pentyneylene. One or more of the hydrogen atoms of each of these groups may be substituted by a like number of halogen atoms such as fluorine, chlorine and/or bromine atoms. Among the above groups, trimethylene, tetramethylene and pentamethylene are particularly preferred.


20 In the ring P, Y is a group



- 13 -

wherein B, D, E₁, E₂, W and m have the same meanings as defined above. The group, which is contained in the above group and is represented by the following formula:

5




wherein E₁, E₂ and W have the same meanings as defined above, is a heterocyclic group derived from piperidine or piperazine, and two or less of the hydrogen atoms on the ring may be substituted by a like number of alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl.

When the above group is a heterocyclic group derived from piperidine, m stands for 0 or 1 (with the proviso that m stands for 1 when Y represents C=), and B represents an oxygen atom, a sulfur atom, a carbonyl group, a sulfinyl group, a sulfonyl group, an alkylene group (an alkylene group having 1-4 carbon atoms preferably, with a methylene group being particularly preferred), an alkenylene group (an alkenylene group having 2-5 carbon atoms preferably, with a 2-propenylene group being particularly preferred), a group -C(OH)R₁- in which R₁ represents an aryl group


- 14 -

having 6-14 carbon atoms, such as phenyl or naphthyl, and one or more hydrogen atoms may be substituted, a group $-\text{CHR}_2-$ in which R_2 represents an aryl group having 6-14 carbon atoms, such as phenyl or naphthyl, and
 5 one or more hydrogen atoms may be substituted, a group



in which the double bond is coupled with W, R_3 represents an aryl group having 6-14 carbon atoms, such as phenyl or naphthyl, or an aralkyl group having 7-22
 10 carbon atoms, such as benzyl or phenethyl, and these groups may be in substituted forms, or a cyclic or acyclic acetal group in which one or more of the hydrogen atoms may be substituted.

Examples of the cyclic or acyclic acetal group
 15 can include:



In the above-described definition of B, preferred examples of the substituents for the groups R_1, R_2 can include alkyl groups having 1-4 carbon atoms preferably, such as methyl and ethyl; and aryl groups having 6-
 20 14 carbon atoms, such as phenyl and naphthyl. These groups may be substituted by one or more of halogen

- 15 -

atoms such as fluorine, chlorine and/or bromine, alkoxy groups having 1-4 carbon atoms preferably, such as methoxy and/or ethoxy, hydroxyl groups, cyano groups, nitro groups and the like.

5 Exemplary substituents for R_3 can include one or more of halogen atoms such as fluorine, chlorine and/or bromine, alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl, alkoxy groups having 1-4 carbon atoms preferably, such as methoxy and/or
10 ethoxy, and hydroxyl groups. Illustrative of the substituent for the cyclic or acyclic acetal can be halogen atoms such as fluorine, chlorine and bromine, alkyl groups having 1-4 carbon atoms preferably, such as methyl and ethyl, aryl groups having 6-14 carbon
15 atoms, such as phenyl and naphthyl, aralkyl groups having 7-22 carbon atoms, such as benzyl and phenethyl, and alkylidene groups having 1-4 carbon atoms preferably, such as methylenidene and ethylenidene.

Among these examples of B, particularly preferred
20 is a carbonyl group.

When the heterocyclic group is a group derived from piperazine, m stands for 0 or 1 (preferably 0), and B represents a carbonyl group, a sulfonyl group, an alkylene group (preferably, an alkylene group having 1-
25 4 carbon atoms, with a methylene group being particu-

- 16 -

larly preferred), an alkenylene group (preferably, an alkenylene group having 3-6 carbon atoms, with a 2-propenylene group being particularly preferred), a group -CHR₄- in which R₄ represents an aryl group having 6-14 carbon atoms, such as phenyl or naphthyl.

The above-described R₄ may be substituted further by one or more of halogen atoms such as fluorine, chlorine and/or bromine, alkyl groups having 1-4 carbon atoms preferably, such as methyl and/or ethyl, and/or alkoxy groups having 1-4 carbon atoms preferably, such as methoxy and/or ethoxy.

Among the above-described examples of B, preferred is a substituted or unsubstituted phenyl-methylene group.

Preferred examples of D can include aromatic hydrocarbon groups having 6-28 carbon atoms preferably, such as a phenyl group in which one or more of the hydrogen atoms may be substituted and a naphthyl group in which one or more of the hydrogen atoms may be substituted.

Other preferred examples of D can include aromatic heterocyclic groups, preferably those each of which is monocyclic or dicyclic and contains three or less hetero atoms, such as pyridyl, pyrimidinyl, benzisothiazolyl, benzisoxazolyl, indazolyl and indolyl

- 17 -

groups in which one or more of hydrogen atoms may be substituted. Illustrative of the hetero atoms can be oxygen, sulfur and/or nitrogen atoms.

Examples of the substituents for the above
5 aromatic hydrocarbon group or aromatic heterocyclic group can include halogen atoms such as fluorine, chlorine and bromine; alkyl groups having 1-4 carbon atoms preferably, such as methyl and ethyl; alkoxy groups having 1-4 carbon atoms preferably, such as
10 methoxy and ethoxy; aryl groups having 6-14 carbon atoms, such as phenyl and naphthyl; aralkyl groups having 7-22 carbon atoms, such as benzyl and phenethyl; aralkyloxy groups having 7-22 carbon atoms preferably, such as benzyloxy; cyano groups; nitro groups; carboxyl
15 groups; alkoxycarbonyl groups (with an alcohol moiety thereof having 1-6 carbon atoms preferably); lower alkylsulfonylamino groups (with an alkyl moiety thereof having 1-4 carbon atoms preferably); carbamoyl groups; and hydroxyl groups.

20 Among these examples of group D, preferred ones can include phenyl groups which may be unsubstituted or substituted by one or more of halogen atoms, alkoxy groups and/or hydroxyl groups; benzisothiazolyl groups which may be unsubstituted or substituted by one or
25 more halogen atoms; benzisoxazolyl groups which may be

- 18 -

unsubstituted or substituted by one or more halogen atoms; and indazolyl groups which may be unsubstituted or substituted by one or more halogen atoms. Particularly preferred are an unsubstituted phenyl group; and
5 phenyl groups substituted by one or more of fluorine atoms, methoxy groups and/or hydroxyl groups.

Many of the compounds (I) according to the present invention have isomers. It is to be noted that these isomers and mixtures thereof are all embraced by
10 the present invention.

Various processes can be employed for the preparation of the pyrroloazepine compounds (I) according to the present invention. It is however preferred to prepare them, for example, by any one of the processes
15 to be described as Process 2 onwards while using as starting material pyrroloazepine compounds (II) or (II') available by Process 1 which will be described hereinafter.


Process 1:

20 The pyrroloazepine compounds (II) and (II') useful as starting materials can be synthesized, for example, by the following process:

Each compound of the formula (II) or the formula (II') can be obtained in accordance with the following
25 reaction scheme, namely, by reacting a pyrrole-3-

- 19 -

carboxylic acid or a derivative thereof represented by the formula (XV) with a β -aminopropionic acid or a derivative thereof represented by the formula (XVI') or an organic or inorganic salt thereof and, if necessary, conducting deprotection to obtain a compound represented by the formula (XVII') and then subjecting the thus-obtained compound to a ring-closing reaction.



wherein Q represents a hydroxyl group, an alkoxy group or an eliminative group easily replaceable by an amino group, and R and R_7 have the same meanings as defined above.

Examples of the eliminative group, which is easily replaceable with an amino group and is represented by the group Q in the compound (XV), can include halogen atoms, an acyloxy group and a p-nitrophenoxy

- 20 -

group. On the other hand, as the carboxyl-protecting group represented by the group R₇ in the compound (XVI'), it is possible to use, in addition to lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl and aralkyl groups having 7-20 carbon atoms, such as benzyl and 9-anthrylmethyl, conventional protecting groups such as those described in T.W. Greene: "Protective Groups in Organic Synthesis" (John Wiley & Sons, Inc.) and the like.

For the synthesis of the compound (XVII'), it is possible to use any one of various processes disclosed in "Compendium of Organic Synthetic Methods" (WILEY-INTERSCIENCE; A Division of John Wiley & Sons, Inc.) and the like.

Illustrative synthesis processes of the compound (XVII') can include a process in which a pyrrole-3-carboxylic acid [the compound (XV) in which Q=OH] and a β -aminopropionic acid or a derivative thereof represented by the formula (XVI') or an organic or inorganic salt thereof are treated with an organic compound such as diethyl phosphorocyanidate (DEPC), diphenylphosphoryl azide (DPPA), dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride or 2-iodo-1-methylpyridinium

- 21 -

iodide or an inorganic compound such as silicon tetrachloride or tin tetrachloride, if necessary, in the presence of an organic or inorganic base; and a process in which a pyrrole-3-carboxylic acid is converted into
5 its acid halide, symmetric acid anhydride, mixed acid anhydride, its active ester such as p-nitrophenyl ester, or the like by a method known *per se* in the art, and is then reacted with the compound (XVI'), if necessary, in the presence of an organic or inorganic base.

10 Each compound (XVII') thus obtained is subjected to a cyclization reaction, optionally after removing the protecting group by virtue of a suitable method such as the action of an acid or a base, or catalytic reduction. This cyclization reaction is conducted by
15 treating the compound (XVII') together with an organic acid such as methanesulfonic acid, an inorganic acid such as sulfuric acid or polyphosphoric acid or a mixture of such an organic or inorganic acid and phosphorus pentoxide at room temperature to 170°C,
20 preferably at 80-120°C. In this case, a solvent which does not take part in the reaction may be added as needed.

As an alternative, the cyclization reaction can also be practiced by, optionally after addition of a
25 catalyst, treating the compound (XVII') with oxalyl

- 22 -

chloride, thionyl chloride, thionyl bromide, oxalyl
bromide, phosgene, phosphorus trichloride, phosphorus
tribromide, phosphoryl chloride, phosphoryl bromide or
the like to convert it into its corresponding acid
5 halide and then treating the acid halide at -20°C to
reflux temperature in the presence of a Lewis acid such
as aluminum chloride, aluminum bromide, boron
trifluoride-ether complex or tin tetrachloride in a
solvent such as dichloromethane, 1,2-dichloroethane or
10 nitromethane. In the above-described reactions, the
compound (II) and the compound (II') can be formed at
varied ratios by changing the reaction conditions.

Process 2:


Among the pyrroloazepine compounds (I), compounds
15 (Ia) and (Ia') in each of which Z_1 and Z_2 are combined
together to represent an oxygen atom can be synthesized,
for example, by any one of the following processes.

Process (a)

Each compound (Ia) or compound (Ia') can be ob-
20 tained in accordance with the following reaction
scheme, namely, by reacting a compound represented by
the formula (II) or (II') with a compound represented
by the formula (III) to convert the compound (II) or
(II') into a compound represented by the formula (IV)
25 or (IV') and then reacting a nitrogen-containing com-

- 23 -

pound represented by the formula (V) or a salt thereof to the compound (IV) or (IV').



5 wherein A, R, X, X' and Y have the same meanings as defined above.

In the above-described reactions, the conversion from the compound (II) or (II') into the compound (IV) or (IV') can be effected by treating the compound (II) or (II') with an organic or inorganic base and then reacting the compound (III), or by causing the compound (III) to act on the compound (II) or (II') in the presence of such a base.

15 The groups X and X' in the compound (III) are eliminative groups. Illustrative can be halogen atoms such as chlorine and bromine, alkylsulfonyloxy groups such as methanesulfonyloxy, and arylsulfonyloxy groups

- 24 -

such as p-toluenesulfonyloxy.

Exemplary organic or inorganic bases can include sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, triethylamine, sodium ethoxide, and potassium t-butoxide. Further, illustrative solvents usable in the above reaction can include acetone, 2-butanone, acetonitrile, dimethyl sulfoxide, dioxane and toluene. The reaction can be conducted at -20°C to reflux temperature.

To prepare the compound (Ia) or (Ia') from the thus-obtained compound (IV) or (IV'), it is only necessary to react the compound (IV) or (IV') with the nitrogen-containing compound (V) or an organic acid salt or inorganic acid salt thereof in a solventless manner or in a solvent such as the above-described solvent, methanol or ethanol at room temperature to 150°C. In this reaction, an organic base such as triethylamine, pyridine, collidine or potassium t-butoxide or an inorganic base such as potassium carbonate, sodium carbonate, sodium hydrogencarbonate, sodium hydroxide or potassium hydroxide can be used as needed. Further, an alkali iodide such as potassium iodide or sodium iodide can also be added as needed.

Examples of the nitrogen-containing compound (V)

- 25 -


can include 1-phenylpiperazine, 1-(2-fluorophenyl)-
piperazine, 1-(3-fluorophenyl)piperazine, 1-(4-fluoro-
phenyl)piperazine, 1-(4-hydroxyphenyl)piperazine, 1-(2-
chlorophenyl)piperazine, 1-(3-chlorophenyl)piperazine,
5 1-(4-chlorophenyl)piperazine, 1-(2-methoxyphenyl)-
piperazine, 1-(3-methoxyphenyl)piperazine, 1-(4-
methoxyphenyl)piperazine, 1-(4-methanesulfonamido-
phenyl)piperazine, 1-(4-cyanophenyl)piperazine, 1-(4-
carbamoylphenyl)piperazine, 1-(4-methoxycarbonyl-
10 phenyl)piperazine, 1-(2-pyridyl)piperazine, 1-(2-
pyrimidinyl)piperazine, 1-benzylpiperazine, 1-diphenyl-
methylpiperazine, 1-cinnamylpiperazine, 1-benzoyl-
piperazine, 1-(4-benzyloxybenzoyl)piperazine, 1-(4-
hydroxybenzoyl)piperazine, 1-(2-furoyl)piperazine, 1-
15 (1,2-benzisoxazol-3-yl)piperazine, 1-(1,2-benz-
isothiazol-3-yl)piperazine, 4-phenylpiperidine, 4-
benzylpiperidine, α,α -bis(4-fluorophenyl)-4-piperidine-
methanol, 4-(4-fluorobenzoyl)piperidine, 4-benzoyl-
piperidine, 4-(4-methoxybenzoyl)piperidine, 4-(4-
20 chlorobenzoyl)piperidine, 3-(4-fluorobenzoyl)-
piperidine, 4-(6-fluoro-1,2-benzisoxazol-3-yl)-
piperidine, 4-(6-fluoro-1,2-benzisothiazol-3-yl)-
piperidine, 4-(6-fluoro-1H-indazol-3-yl)piperidine, 4-
(4-fluorophenoxy)piperidine, 4-[(4-fluorophenyl)thio]-
25 piperidine, 4-[(4-fluorophenyl)sulfinyl]piperidine, 4-

- 26 -

[(4-fluorophenyl)sulfonyl]piperidine, 4-[bis(4-fluorophenyl)methylene]piperidine, and 4-(4-fluorobenzoyl)-piperidine ethylene acetal. They are either known or can be readily prepared by known processes or processes similar to such known processes.

Process (b)

Further, the compound (Ia) or (Ia') can be obtained by causing a compound represented by the formula (VI) to act on the compound represented by the formula (II) or (II') in accordance with the following reaction formula:



wherein A, R, X and Y have the same meanings as defined above.

- 27 -

The conversion from the compound (II) or (II') into the compound (Ia) or (Ia') is conducted by causing the compound (VI) to act either after treatment of the compound (II) or (II') with an inorganic base or an organic base or in the presence of an inorganic base or an organic base. Reaction conditions are similar to those employed upon conversion from the compound (II) into the compound (IV) and described above under Process (a) of Process 2. Further, the compound (VI) can be synthesized by reacting the compound (III) with the compound (V) in a manner known *per se* in the art.

Process 3:

Among the pyrroloazepine compounds (I), the compounds (Ic), (Ic'), (If) and (If') in each of which Z_1 and Z_2 both represent groups SR_5 (in which R_5 has the same meaning as described above) or Z_1 and Z_2 are combined together to represent the group




wherein G has the same meaning as defined above can be synthesized by any one of the following processes.

Process (a)

The compound (If) or (If') is obtained in accordance with the following reaction scheme, namely, by

- 28 -

reacting a thiol compound, which is represented by the formula (VIIa) or (VIIb) [the compound (VIIa) and the compound (VIIb) may hereinafter be collectively called "the thiol compound (VII)"], with a compound (II) or (II') and then causing a compound (VI) to act.



wherein A, G, R, R₅, X, Y, Z₁' and Z₂' have the same meanings as defined above.


For the conversion from the compound (II) or (II') into the compound (IX) or (IX'), a suitable method can be selected from those disclosed, for example, in T.W. Greene: "Protective Groups in Organic Synthesis" (John Wiley & Sons, Inc.) and the like. Describing one example, there is a process in which the thiol compound (VII) and boron trifluoride-ether complex are caused to act on the compound (II) or (II') in

- 29 -

chloroform. Further, the conversion from the compound (IX) or (IX') into the compound (If) or (If') can be effected under the same conditions as in the conversion from the compound (II) into the compound (Ia) described above under Process (b) of Process 2.

Process (b)

Each compound represented by the formula (Ic) or (Ic') can be obtained by causing the thiol compound (VII) to act on a compound (Ib) or (Ib') in accordance with the following reaction scheme.



wherein A, G, R, R₅, Y', Z₁' and Z₂' have the same meanings as described above.

The conversion from the compound (Ib) or (Ib')

- 30 -


into the compound (Ic) or (Ic') can be effected under similar conditions as in the conversion of from the compound (II) into the compound (IX) described above under Process (a) of Process 3.

5 Process 4:


Among the pyrroloazepine compounds (I), the compounds (Id) and (Id') and the compounds (Ig) and (Ig') in each of which Z_1 and Z_2 are combined together to represent a group NOR_6 can each be synthesized by any
 10 one of the following processes.

Process (a)

Each compound (Ig) or (Ig') is obtained in accordance with the following reaction scheme, namely, by causing hydroxylamine or a derivative thereof (VIII) or
 15 a salt thereof to act on a compound represented by the formula (IV) or (IV') and then causing a nitrogen-containing compound (V) to act.



- 31 -



wherein A, R, R₆, X and Y have the same meanings as defined above.

The reaction between the compound (IV) or (IV') and the hydroxylamine or its derivative (VIII) is effected, if necessary, in the presence of an organic base such as pyridine, triethylamine, collidine or sodium acetate or an inorganic base such as potassium carbonate or sodium hydroxide. The hydroxylamine or its derivative (VIII) may also be used in the form of an organic acid salt or an inorganic acid salt.


The reaction is conducted at 0°C to reflux temperature, preferably 0°C-100°C by adding a suitable solvent, for example, methanol, ethanol, propanol, tetrahydrofuran, dimethylformamide or dimethyl sulfoxide as needed.

Further, the conversion from the thus-obtained compound (X) or (X') into the compound (Ig) or (Ig') can be effected under similar conditions as in the conversion from the compound (IV) into the compound (Ia) shown above under Process (a) of Process 2.

- 32 -

Process (b)

Each compound (Id) or (Id') is obtained by causing hydroxylamine or its derivative (VIII) or a salt thereof to act on a compound (Ib) or (Ib') in accordance with the following reaction formula.



wherein A, R, R_6 and Y have the same meanings as defined above.

The conversion from the compound (Ib) or (Ib') into the compound (Id) or (Id') can be effected under similar conditions as the conversion from the compound (IV) into the compound (X) shown above under Process (a) of Process 4.


- 33 -

Process 5:

Among the pyrroloazepine compounds (I), the compounds (Ie) and (Ie') and the compounds (Ih) and (Ih') in each of which Z_1 represents a hydrogen atom and Z_2 represents a hydroxyl group can each be synthesized by

Process (a)

Each compound (Ih) or (Ih') is obtained in accordance with the following reaction scheme, namely, by reducing a compound represented by the formula (IV) or (IV') and then causing a nitrogen-containing compound (V) to act.



wherein A, R, X and Y have the same meanings as defined above.


The conversion from the compound (IV) or (IV')

into the compound (XI) or (XI') is conducted by treating the compound (IV) or (IV') with a reducing agent such as sodium borohydride, potassium borohydride, sodium cyanoborohydride or tri(n-butyl)tin hydride at -78°C to reflux temperature, preferably -20°C to room temperature in an ordinarily-employed solvent.


The conversion from the compound (XI) or (XI') into the compound (Ih) or (Ih') can be effected under similar conditions as the conversion from the compound (IV) into the compound (Ia) shown above under Process (a) of Process 2.

Process (b)

Each compound (Ie) or (Ie') is obtained by reducing a compound represented by the formula (Ib) or (Ib') in accordance with the following reaction formula.



- 35 -



wherein A, R and Y' have the same meanings as defined above.

The conversion from the compound (Ib) or (Ib') into the compound (Ie) or (Ie') can be effected under similar conditions as in the conversion from the compound (IV) into the compound (XI) shown above under Process (a) of Process 5.

Process 6:


Among the pyrroloazepine compounds (I), the compounds (Ii) or (Ii') in each of which the bond indicated by the dashed line is present and Z₁ represents a hydrogen atom can be synthesized by any one of the following processes.

Process (a)

Each compound (Ii) or (Ii') is obtained in accordance with the following reaction scheme, namely, by subjecting a compound represented by the formula (XI) or (XI') to a dehydrating reaction to obtain a compound represented by the formula (XII) or (XII') and then

- 36 -

causing a nitrogen-containing compound (V) to act on the compound (XII) or (XII').



5 wherein A, R, X and Y have the same meanings as defined above.

In the above-described reaction, the conversion from the compound (XI) or (XI') into the compound (XII) or (XII') can be effected by treating the compound (XI) or (XI') with an acid such as hydrogen chloride,
 10 hydrogen bromide, sulfuric acid, methanesulfonic acid or p-toluenesulfonic acid at -20°C to 100°C, preferably at -20°C to room temperature in a solvent such as water, methanol, ethanol, ethyl acetate, chloroform or
 15 toluene.

As an alternative, the conversion into the compound (XII) or (XII') can also be effected by causing


- 37 -

methanesulfonyl chloride, p-toluenesulfonyl chloride, phosphorus trichloride, phosphorus oxychloride, thionyl chloride or the like and a base such as triethylamine, pyridine or collidine to act on the compound (XI) or
5 (XI') in a solvent such as dichloromethane, chloroform or toluene.


The conversion from the compound (XII) or (XII') into the compound (Ii) or (Ii') can be effected under similar conditions as in the conversion from the compound (IV) into the compound (Ia) described above under
10 Process (a) of Process 2.

Process (b)

Each compound (Ii) or (Ii') is also obtained by subjecting a compound represented by the formula (Ih) or (Ih') to a dehydrating reaction in accordance with
15 the following reaction formula:



- 38 -




wherein A, R and Y have the same meanings as defined above.

In the above-described reaction, the conversion
 5 from the compound (Ih) or (Ih') into the compound (Ii)
 or (Ii') can be effected under similar conditions as in
 the conversion from the compound (XI) into the compound
 (XII) described above under Process (a) of Process 6.

Process 7:

10 Among the pyrroloazepine compounds (I), compounds
 (Ij) or (Ij') in each of which Z_1 and Z_2 both represent
 hydrogen atoms can be obtained in accordance with the
 following reaction scheme, namely, by reducing a com-
 pound represented by the formula (XII) or (XII') to ob-
 15 tain a compound represented by the formula (XIII) or
 (XIII') and then reacting a nitrogen-containing com-
 pound (V) to the compound (XIII) or (XIII').

- 39 -



wherein A, R, X and Y have the same meanings as defined above.

5 In the above-described reaction, the conversion from the compound (XII) or (XII') into the compound (XIII) or (XIII') can be conducted by treating, in the presence of a catalyst such as palladium-carbon or platinum, the compound (XII) or (XII') with hydrogen
10 gas in an ordinarily-employed solvent at -78°C to reflux temperature, preferably at room temperature. Further, the conversion from the compound (XIII) or (XIII') into the compound (Ij) or (Ij') can be effected under similar conditions as in the conversion from the
15 compound (IV) into the compound (Ia) described above under Process (a) of Process 2.

 If necessary, the compounds (I) of the present invention obtained according to the above-described

- 40 -

processes can each be reacted with one of various acids to convert the compound into its salt. Then, the resulting salt can be purified by a method such as recrystallization or column chromatography.

5 Exemplary acids usable for the conversion of the pyrroloazepine compounds (I) into their salts can include inorganic acids such as hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid and hydrobromic acid; and organic acids such as maleic acid, fumaric
10 acid, tartaric acid, lactic acid, citric acid, acetic acid, methanesulfonic acid, p-toluenesulfonic acid, adipic acid, palmitic acid and tannic acid.

Further, the compounds (I) according to the present invention include many compounds containing asym-
15 metric centers. Each racemic mixture can be resolved by one or more of various methods, whereby a single enantiomer can be obtained.

Usable methods include, for example:

- (1) Resolution by an optical resolution column.
- 20 (2) Resolution by recrystallization subsequent to conversion into a salt with a chiral acid.
- (3) Resolution by an enzyme reaction.
- (4) Resolution by a combination of the above
methods (1) to (3).

25 The pyrroloazepine compounds (I) and their salts,

- 41 -

which are obtained as described above, have strong serotonin-2 blocking action and also α_1 blocking action as will be demonstrated in tests to be described subsequently herein. From the results of a toxicity test, they have also been found to possess high safety. The compounds according to the present invention can therefore be used as pharmaceuticals for the treatment of circulatory diseases such as ischemic heart diseases, cerebrovascular disturbances, peripheral circulatory disturbances and hypertension.

When the pyrroloazepine compounds (I) according to this invention are used as pharmaceuticals, they can be administered in effective doses as they are. As an alternative, they can also be formulated into various preparation forms by known methods and then administered.

Exemplary preparation forms as pharmaceuticals include orally administrable preparation forms such as tablets, capsules and syrups as well as parenterally administrable preparation forms such as injections and suppositories. Whichever preparation form is used, a known liquid or solid extender or carrier usable for the formulation of the preparation form can be employed.

Examples of such extender or carrier include

- 42 -

polyvinylpyrrolidone, arabic gum, gelatin, sorbit,
cyclodextrin, tragacanth gum, magnesium stearate, talc,
polyethylene glycol, polyvinyl alcohol, silica, lac-
tose, crystalline cellulose, sugar, starch, calcium
5 phosphate, vegetable oil, carboxymethylcellulose,
sodium laurylsulfate, water, ethanol, glycerin, man-
nitol, syrup, and the like.

When the compounds (I) according to the present
invention are used as pharmaceuticals, their dose
10 varies depending on the administration purpose, the
age, body weight, conditions, etc. of the patient to be
administered. In oral administration, the daily dose
may generally be about 0.01-1,000 mg.

The present invention will next be described in
15 further detail by the following examples and tests. It
is however to be noted that the present invention is by
no means limited to the following examples and tests.

Example 1

Synthesis of benzyl 3-(3-pyrrolicarboxamido)-
20 propionate (Compound No. 1)

Into a solution of 1.67 g (15 mmol) of 3-pyrrole-
carboxylic acid and 6.33 g (18 mmol) of β -alanine ben-
zyl ester p-toluenesulfonate in 20 ml of dimethyl-
formamide, a solution of 2.94 g (18 mmol) of diethyl
25 phosphorocyanidate in 10 ml of dimethylformamide was

- 43 -

added dropwise under ice cooling and stirring. A solution of 3.64 g (36 mmol) of triethylamine in 20 ml of dimethylformamide was then added dropwise. The reaction mixture was stirred at 0°C for 1 hour and then at room temperature for 24 hours.

The reaction mixture was concentrated under reduced pressure, followed by the addition of a 3:1 mixed solvent of ethyl acetate and benzene to the residue. The resultant solution was washed successively with a half-saturated aqueous solution of potassium carbonate, water, a 10% aqueous solution of citric acid, water and a saturated aqueous solution of sodium chloride. The solution was dried over anhydrous sodium sulfate and was then concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column ("Merck No. 9385" was used as its silica gel; the same silica gel was employed in the subsequent examples unless otherwise specifically indicated) (eluent: ethyl acetate/hexane = 2:1), whereby 3.62 g of the title compound were obtained (yield: 89%).

Example 2

Synthesis of ethyl N-methyl-3-(3-pyrrole-carboxamido)propionate (Compound No. 2)

Into a suspension of 3.30 g (30 mmol) of 3-

- 44 -

pyrrolicarboxylic acid in 10 ml of dichloromethane, a solution of 5.90 g (45 mmol) of ethyl N-methyl-3-aminopropionate in 50 ml of dichloromethane, a solution of 4.55 g (45 mmol) of triethylamine in 50 ml of dichloromethane, and 8.63 g (45 mmol) of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride were successively added, followed by stirring at room temperature for 3 hours. Dichloromethane was added to the reaction mixture, and the resulting organic layer was washed successively with a 10% aqueous solution of citric acid, a saturated aqueous solution of sodium hydrogencarbonate and a saturated aqueous solution of sodium chloride. The organic layer was dried over anhydrous magnesium sulfate and was then concentrated under reduced pressure, whereby 5.31 g of the title compound were obtained (yield: 79%).

Example 3

Synthesis of 3-(3-pyrrolicarboxamido)propionic acid (Compound No. 3)

Into a solution of 8.29 g (30.4 mmol) of Compound No. 1 in 200 ml of tetrahydrofuran, 829 mg of 5%-palladium-carbon were added, followed by stirring for 63 hours under a hydrogen gas stream. The reaction mixture was filtered and the thus-obtained solid matter was washed with methanol. The filtrate and the washing

- 45 -

were combined and concentrated under reduced pressure. The residue was recrystallized from methanol-diisopropyl ether, whereby 3.89 g of the title compound were obtained (yield: 70%).

5 **Example 4**

Synthesis of N-methyl-3-(3-pyrrolicarboxamido)-propionic acid (Compound No. 4)

A 2 N aqueous solution of sodium hydroxide (58 ml, 116 mmol) was added to 5.16 g (23 mmol) of
10 Compound No. 2, followed by stirring at room temperature for 1.5 hours. After the reaction mixture was washed twice with ethyl ether, 6 N hydrochloric acid was added under ice cooling to adjust the pH of the mixture to 2. The mixture was saturated with sodium
15 chloride. The resultant mixture was extracted four times with ethyl acetate. The extracts were combined, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The thus-obtained crude crystals were recrystallized from ethyl acetate-
20 hexane, whereby 4.10 g of the title compound were obtained (yield: 91%).

Example 5

Synthesis of 1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione (Compound No. 5) and
25 2,4,5,6,7,8-hexahydropyrrolo[3,4-c]azepine-4,8-dione

- 46 -

(Compound No. 6)

A mixture of 310 mg (1.7 mmol) of Compound No. 3 and 15.5 g of polyphosphoric acid (80%) was stirred for 1 hour in an oil bath of 100°C. The reaction mixture
5 was poured into 100 g of ice water, to which a 4 N aqueous solution of sodium hydroxide was added to adjust its pH to 6. The mixture was saturated with sodium chloride, followed by extraction (4 times) with tetrahydrofuran. The organic layer was washed with a
10 saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was fractionated and purified by chromatography on a silica gel (eluent: methanol/ chloroform = 1/19 → 1/9), whereby 166 mg
15 of Compound No. 5 and 73 mg of Compound No. 6 were obtained (yields: 59% and 26%).

Example 6

Synthesis of 5-methyl-1,4,5,6,7,8-hexahydropyrrolo-
[3,2-c]azepine-4,8-dione (Compound No. 7) and 5-
20 methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]azepine-4,8-dione (Compound No. 8)

A mixture of 981 mg (5 mmol) of Compound No. 4 and 50 g of polyphosphoric acid (80%) was stirred for 30 minutes in an oil bath of 100°C. The reaction mixture
25 was poured into 200 g of ice water, to which a 4 N

- 47 -

aqueous solution of sodium hydroxide was added to adjust its pH to 5. The mixture was saturated with sodium chloride, followed by extraction (thrice) with tetrahydrofuran. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was fractionated and purified by chromatography on a silica gel (eluent: methanol/chloroform = 3/97), whereby 653 mg of Compound No. 7 and 178 mg of Compound No. 8 were obtained (yields: 73% and 20%).

Example 7

Synthesis of 1-(4-chlorobutyl)-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione (Compound No. 9)

A suspension of 3.03 g (17 mmol) of Compound No. 7, 9.40 g (68 mmol) of potassium carbonate and 11.66 g (68 mmol) of 1-bromo-4-chlorobutane in 100 ml of 2-butanone was refluxed for 11 hours. The reaction mixture was filtered, and the thus-obtained solid matter was washed with 2-butanone. The filtrate and the washing were combined and concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/99), whereby 4.46 g of the title compound were obtained

- 48 -

(yield: 98%).

Example 8

Synthesis of 1-(4-bromobutyl)-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione (Compound No. 10)

A suspension of 178 mg (1 mmol) of Compound No. 7, 691 mg (5 mmol) of potassium carbonate and 0.6 ml (5 mmol) of 1,4-dibromobutane in 20 ml of dimethylformamide was stirred for 1 hour in an oil bath of 80°C. The reaction mixture was concentrated under reduced pressure. Ice water was added to the residue, followed by neutralization with 6 N hydrochloric acid. The thus-obtained mixture was extracted with chloroform (thrice). The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column (eluent: ethyl acetate), whereby 254 mg of the title compound were obtained (yield: 81%).

Example 9

Synthesis of 2-(4-chlorobutyl)-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]azepine-4,8-dione (Compound No. 11)

A suspension of 891 mg (5 mmol) of Compound No.

- 49 -

8, 2.76 g (20 mmol) of potassium carbonate and 3.43 g (20 mmol) of 1-bromo-4-chlorobutane in 40 ml of 2-butanone was refluxed for 11 hours. The reaction mixture was post-treated in a similar manner as in Example 7 and the residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/49), whereby 1.29 g of the title compound were obtained (yield: 96%).

Example 10

Synthesis of 1-(4-chlorobutyl)-8-hydroxyimino-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepin-4-one (Compound No. 12)

A solution of 134 mg (0.5 mmol) of Compound No. 9 and 42 mg (0.6 mmol) of hydroxylamine hydrochloride in 10 ml of methanol was refluxed for 190 hours. The reaction mixture was concentrated under reduced pressure. A half-saturated aqueous solution of potassium carbonate was added to the residue, followed by extraction (twice) with chloroform. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/99 → 1/49), whereby 70 mg of the title compound were obtained (yield: 49%).

- 50 -

Example 11

Synthesis of 2-(4-chlorobutyl)-8-hydroxyimino-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]azepin-4-one (Compound No. 13)

5 A solution of 537 mg (2 mmol) of Compound No. 11, 278 mg (4 mmol) of hydroxylamine hydrochloride and 328 mg (4 mmol) of sodium acetate in 20 ml of methanol was refluxed for 4 hours. The reaction mixture was post-treated in a similar manner as in Example 10 and the
10 residue was purified by chromatography on a silica gel column (eluent: ethyl acetate), whereby 273 mg of the title compound were obtained (yield: 48%).

Example 12

 Synthesis of 1-(4-chlorobutyl)-8-hydroxy-5-methyl-
15 1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepin-4-one
 (Compound No. 14)

 Into a solution of 269 mg (1 mmol) of Compound No. 9 in 15 ml of ethanol, 378 mg (10 mmol) of sodium borohydride were added under ice cooling and stirring,
20 followed by stirring at room temperature for 18 hours. Water (15 ml) was added to the reaction mixture. The resultant mixture was stirred at room temperature for 2 hours and then concentrated under reduced pressure. Water was added to the residue, followed by extraction
25 (twice) with ethyl acetate. The organic layer was

- 51 -

washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column
5 (eluent: methanol/chloroform = 1/49), whereby 246 mg of the title compound were obtained (yield: 91%).

Example 13

Synthesis of 1-[4-[4-(4-fluorobenzoyl)piperidino]-
butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-
10 c]azepine-4,8-dione (Compound No. 15)

A suspension of 100 mg (0.32 mmol) of Compound No. 10, 364 mg (0.96 mmol) of 4-(4-fluorobenzoyl)-piperidine p-toluenesulfonate, 265 mg (1.92 mmol) of potassium carbonate and 54 mg (0.38 mmol) of sodium
15 iodide in 20 ml of dimethylformamide was stirred for 15 hours in an oil bath of 80°C. Water was added to the reaction mixture, followed by extraction with a 3:1 mixed solvent of ethyl acetate and benzene. The organic layer was washed with water (twice) and a
20 saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/19), whereby 123 mg of the
25 title compound were obtained (yield: 88%).

- 52 -

Example 14

Synthesis of 5-methyl-1-[4-(4-phenylpiperazin-1-yl)butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione (Compound No. 16)

5 A suspension of 269 mg (1 mmol) of Compound No. 9, 162 mg (1 mmol) of N-phenylpiperazine, 168 mg (2 mmol) of sodium hydrogencarbonate and 300 mg (2 mmol) of sodium iodide in 15 ml of acetonitrile was refluxed for 11 hours. The reaction mixture was post-treated in
10 a similar manner as in Example 10 and the residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/49), whereby 360 mg of the title compound were obtained (yield: 91%).

Example 15

15 Synthesis of 1-[4-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidino]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione (Compound No. 17)

 A suspension of 269 mg (1 mmol) of Compound No. 9, 220 mg (1 mmol) of 4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidine, 168 mg (2 mmol) of sodium hydrogencarbonate and 300 mg (2 mmol) of sodium iodide in 15 ml
20 of acetonitrile was refluxed for 20 hours. The reaction mixture was post-treated in a similar manner as in Example 10. The residue was purified by chromatography
25 on a silica gel column (eluent: methanol/chloroform =

- 53 -

3/97), whereby 406 mg of the title compound were obtained (yield: 90%).

Example 16

Synthesis of 2-[4-[4-(4-fluorobenzoyl)piperidino]-
5 butyl]-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-
c]azepine-4,8-dione (Compound No. 18)

A suspension of 134 mg (0.5 mmol) of Compound No.
11, 122 mg (0.5 mmol) of 4-(4-fluorobenzoyl)piperidine
hydrochloride, 168 mg (2 mmol) of sodium hydrogencar-
10 bonate and 150 mg (1 mmol) of sodium iodide in 10 ml
of acetonitrile was refluxed for 25 hours. The reac-
tion mixture was post-treated in a similar manner as in
Example 10 and the residue was purified by chromato-
graphy on a silica gel column (eluent: methanol/
15 chloroform = 3/97), whereby 186 mg of the title com-
pound were obtained (yield: 85%).

Example 17

Synthesis of 1-[4-[4-(4-fluorobenzoyl)piperidino]-
butyl]-8-hydroxyimino-5-methyl-1,4,5,6,7,8-hexa-
20 hydropyrrolo[3,2-c]azepin-4-one (Compound No. 19)

A suspension of 57 mg (0.2 mmol) of Compound NO.
12, 54 mg (0.22 mmol) of 4-(4-fluorobenzoyl)piperidine
hydrochloride, 74 mg (0.88 mmol) of sodium hydrogencar-
bonate and 66 mg (0.44 mmol) of sodium iodide in 10 ml
25 of acetonitrile was refluxed for 40 hours. The reac-

- 54 -

tion mixture was post-treated in a similar manner as in Example 10 and the residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 3/97 → 1/19), whereby 72 mg of the title compound were obtained (yield: 79%).

Example 18

Synthesis of 8-hydroxyimino-5-methyl-1-[4-(4-phenylpiperazin-1-yl)butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepin-4-one (Compound No. 20)

A solution of 70.4 mg (0.18 mmol) of Compound No. 16 and 29 mg (0.4 mmol) of hydroxylamine hydrochloride in 5 ml of pyridine was stirred for 77 hours in an oil bath of 70°C. The reaction mixture was post-treated in a similar manner as in Example 10 and the residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 3/97), whereby 47 mg of the title compound were obtained (yield: 64%).

Example 19

Synthesis of 2-[4-[4-(4-fluorobenzoyl)piperidino]-butyl]-8-hydroxyimino-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]azepin-4-one (Compound No. 21)

A suspension of 142 mg (0.5 mmol) of Compound No. 13, 122 mg (0.5 mmol) of 4-(4-fluorobenzoyl)piperidine hydrochloride, 168 mg (2 mmol) of sodium hydrogencarbonate and 150 mg (1 mmol) of sodium iodide in 10 ml

- 55 -

of acetonitrile was refluxed for 22 hours. The reaction mixture was post-treated in a similar manner as in Example 10 and the residue was purified by chromatography on a silica gel column (eluent: methanol/
5 chloroform = 1/9), whereby 202 mg of the title compound were obtained (yield: 89%).

Example 20

Synthesis of 1-[4-[4-(4-fluorobenzoyl)piperidino]-
butyl]-8-hydroxy-5-methyl-1,4,5,6,7,8-hexahydro-
10 pyrrolo[3,2-c]azepin-4-one (Compound No. 22)

A suspension of 217 mg (0.8 mmol) of Compound No. 14, 214 mg (0.88 mmol) of 4-(4-fluorobenzoyl)piperidine hydrochloride, 296 mg (3.52 mmol) of sodium hydrogen-
carbonate and 264 mg (1.76 mmol) of sodium iodide in 15
15 ml of acetonitrile was refluxed for 24 hours. The reaction mixture was post-treated in a similar manner as in Example 10 and the residue was purified by chromatography on a silica gel column (eluent: methanol/chloroform = 1/19 → 2/23), whereby 295 mg
20 of the title compound were obtained (yield: 84%).

In the same manner or a similar manner as in any of the Examples described above, the following compounds are obtained.

- 56 -

- (1) 1-[3-[4-(4-Fluorobenzoyl)piperidino]propyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- 5 (2) 1-[5-[4-(4-Fluorobenzoyl)piperidino]pentyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- (3) 2-[3-[4-(4-Fluorobenzoyl)piperidino]propyl]-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]-azepine-4,8-dione
- 10 (4) 2-[5-[4-(4-Fluorobenzoyl)piperidino]pentyl]-5-methyl-2,4,5,6,7,8-hexahydropyrrolo[3,4-c]-azepine-4,8-dione
- (5) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione
- 15 (6) 5-Ethyl-1-[4-[4-(4-fluorobenzoyl)piperidino]-butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- (7) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-5-propyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- 20 (8) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-5-isopropyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione

- 57 -

- (9) 5-Butyl-1-[4-[4-(4-fluorobenzoyl)piperidino]-butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- 5 (10) 5-Benzyl-1-[4-[4-(4-fluorobenzoyl)piperidino]-butyl]-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- (11) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-5-methyl-1,4,5,6-tetrahydropyrrolo[3,2-c]azepin-4-one
- 10 (12) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepin-4-one
- (13) 8,8-Bis(ethylthio)-1-[4-[4-(4-fluorobenzoyl)-piperidino]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepin-4-one
- 15 (14) 1-[4-[4-(4-Fluorobenzoyl)piperidino]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepin-4-one-8-spiro-2'-(1',3'-dithiolane)
- (15) 1-[4-[4-(2-Furoyl)piperazin-1-yl]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-azepine-4,8-dione
- 20 (16) 1-[4-[4-[Bis(4-fluorophenyl)methylene]-piperidino]butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]azepine-4,8-dione

- 58 -

- (17) 1-[4-[4-(6-Fluoro-1H-indazol-3-yl)piperidino]-
butyl]-5-methyl-1,4,5,6,7,8-hexahydropyrrolo-
[3,2-c]azepine-4,8-dione
- 5 (18) 1-[4-[4-(6-Fluoro-1,2-benzisothiazol-3-yl)-
piperidino]butyl]-5-methyl-1,4,5,6,7,8-
hexahydropyrrolo[3,2-c]azepine-4,8-dione
- (19) 1-[4-[4-(4-Chlorobenzoyl)piperidino]butyl]-5-
methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-
azepine-4,8-dione
- 10 (20) 1-[4-[4-(4-Fluorophenoxy)piperidino]butyl]-5-
methyl-1,4,5,6,7,8-hexahydropyrrolo[3,2-c]-
azepine-4,8-dione
- (21) 1-[4-[4-(6-Fluoro-1,2-benzisoxazol-3-yl)-
piperidino]butyl]-1,4,5,6,7,8-hexahydropyrrolo-
15 [3,2-c]azepine-4,8-dione
- (22) 5-Ethyl-1-[4-[4-(6-fluoro-1,2-benzisoxazol-3-
yl)piperidino]butyl]-1,4,5,6,7,8-hexahydro-
pyrrolo[3,2-c]azepine-4,8-dione
- (23) 1-[4-[4-(6-Fluoro-1,2-benzisothiazol-3-yl)-
20 piperidino]butyl]-1,4,5,6,7,8-hexahydropyrrolo-
[3,2-c]azepine-4,8-dione
- (24) 5-Ethyl-1-[4-[4-(6-fluoro-1,2-benzisothiazol-3-
yl)piperidino]butyl]-1,4,5,6,7,8-hexahydro-
pyrrolo[3,2-c]azepine-4,8-dione

- 59 -

(25) 2-[4-[4-(6-Fluoro-1,2-benzisoxazol-3-yl)-
piperidino]butyl]-5-methyl-2,4,5,6,7,8-
hexahydropyrrolo[3,4-c]azepine-4,8-dione

Physical data of the compounds obtained above in





5 Examples 1-20 are shown in Tables 1-6.

Table 1

Comp'd No.	Structural formula	Property (Melting point (recrystallization solvent))	NMR (δ ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
1		Brown oil	(270MHz) 2.64 (2H, t, J=5.9Hz), 3.65 (2H, m), 5.11 (2H, s), 6.33 (1H, m), 6.59 (1H, m), 6.68 (1H, m), 7.22 (1H, m), 7.29-7.31 (5H, m), 9.90 (1H, br. s)	(film) 3271, 2954, 1732, 1634, 1567, 1520, 1338, 1254, 1210, 1172, 756, 698
2		Brown oil	(400MHz) 1.26 (3H, t, J=7.1Hz), 2.67 (2H, t, J=7.1Hz), 3.18 (3H, s), 3.81 (2H, t, J=7.1Hz), 4.15 (2H, q, J=7.1Hz), 6.44 (1H, m), 6.74 (1H, m), 7.18 (1H, m), 8.73 (1H, br. s)	(film) 3232, 2980, 1731, 1594, 1546, 1508, 1438, 1401, 1375, 1314, 1190, 1106, 1046, 968, 756
3		Colorless needle crystals 178.5-183.0°C (methanol- isopropyl ether)	(400MHz) (DMSO-d ₆ /TMS) 2.45 (2H, t, J=7.1Hz), 3.36 (2H, m), 6.43 (1H, s), 6.71 (1H, s), 7.26 (1H, s), 7.72 (1H, m), 11.04 (1H, br. s), 12.09 (1H, s)	(KBr) 3381, 3272, 1718, 1567, 1538, 1427, 1351, 1210, 853, 757
4		Pale yellow powdery crystals 125.0-127.5°C (ethyl acetate- hexane)	(400MHz) (DMSO-d ₆ /TMS) 2.52 (2H, m), 3.05 (3H, s), 3.63 (2H, t, J=7.3Hz), 6.29 (1H, m), 6.74 (1H, m), 7.12 (1H, m), 11.11 (1H, br. s), 12.24 (1H, br. s)	(KBr) 3330, 3122, 1721, 1574, 1516, 1407, 1305, 1211, 1098, 906, 752





* Measured in CDCl₃ with TMS as an internal standard unless otherwise specifically indicated.

Table 2

Comp'd No.	Structural formula	Property Melting point (recrystallization solvent)	NMR (δ ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
5		Colorless needle crystals 285-287°C (decomp'd) (methanol- isopropyl ether)	(400MHz) (DMSO- d_6 /TMS) 2.71 (2H, m), 3.33 (2H, m), 6.57 (1H, s), 7.11 (1H, d, J=2.4Hz), 8.29 (1H, m), 12.13 (1H, br. s)	(KBr) 3306, 3037, 2956, 1642, 1503, 1438, 1407, 1395, 1268, 882, 762
6		Colorless powdery crystals 287°C min. (methanol- isopropyl ether)	(400MHz) (DMSO- d_6 /TMS) 2.65 (2H, m), 3.29 (2H, m), 7.34 (1H, s), 7.43 (1H, s), 7.80 (1H, m), 11.97 (1H, br. s)	(KBr) 3309, 3120, 3057, 2946, 2880, 1647, 1620, 1526, 1474, 1456, 1419, 1377, 1360, 910, 838, 810, 759
7		Colorless prism crystals 224.0-225.5°C (chloroform-hexane)	(400MHz) 2.86 (2H, m), 3.23 (3H, s), 3.71 (2H, m), 6.89 (1H, t, J=2.8Hz), 7.04 (1H, t, J=2.8Hz), 9.41 (1H, br. s)	(KBr) 3186, 1664, 1611, 1548, 1500, 1401, 1365, 1277, 1232, 1188, 1112, 1090, 955, 927, 890, 827, 785, 761, 692
8		Colorless prism crystals 175.0-176.5°C (chloroform-hexane)	(400MHz) 2.84 (2H, m), 3.21 (3H, s), 3.68 (2H, m), 7.41 (1H, m), 7.47 (1H, m), 10.32 (1H, br. s)	(KBr) 3108, 2956, 1622, 1522, 1490, 1451, 1391, 1243, 1171, 1098, 958, 924, 863, 815, 758





* Measured in CDCl_3 with TMS as an internal standard unless otherwise specifically indicated.

Table 3

Comp'd No.	Structural formula	Property Melting point (recrystallization solvent)	NMR (δ , ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
9		Colorless needle crystals 88.0-89.0°C (ethanol-isopropyl ether)	(270MHz) 1.80 (2H, m), 1.91 (2H, m), 2.85 (2H, m), 3.22 (3H, s), 3.53 (2H, t, J=6.6Hz), 3.66 (2H, m), 4.35 (2H, t, J=6.6Hz), 6.79 (1H, d, J=2.6Hz), 6.89 (1H, d, J=2.6Hz)	(KBr) 2937, 1630, 1522, 1504, 1410, 1318, 1246, 1178, 1095, 912, 862, 798, 750, 724
10		Colorless needle crystals 92.0-93.0°C (ethyl acetate-hexane)	(270MHz) 1.80-1.99 (4H, m), 2.85 (2H, m), 3.22 (3H, s), 3.39 (2H, t, J=5.9Hz), 3.66 (2H, m), 4.34 (2H, t, J=6.6Hz), 6.79 (1H, d, J=2.6Hz), 6.89 (1H, d, J=2.6Hz)	(KBr) 3630, 2934, 1649, 1630, 1505, 1410, 1247, 1180, 1098, 911, 752
11		Colorless powdery crystals 112.0-113.5°C (chloroform-ether)	(400MHz) 1.77 (2H, m), 1.99 (2H, m), 2.81 (2H, m), 3.19 (3H, s), 3.54 (2H, t, J=6.3Hz), 3.66 (2H, m), 3.96 (2H, t, J=7.0Hz), 7.33 (2H, m)	(KBr) 3116, 2948, 1655, 1624, 1540, 1519, 1490, 1406, 1318, 1244, 1169, 925, 879, 761
12		Colorless prism crystals 194.0-195.5°C (ethyl acetate-hexane)	(400MHz) (DMSO-d ₆ /TMS) 1.63 (2H, m), 1.78 (2H, m), 2.89 (2H, m), 2.97 (3H, s), 3.42 (2H, m), 3.58 (2H, t, J=6.6Hz), 4.19 (2H, t, J=6.9Hz), 6.36 (1H, d, J= 2.9Hz), 6.94 (1H, d, J= 2.9Hz)	(KBr) 3154, 2937, 1591, 1508, 1397, 1322, 1250, 1203, 1099, 1028, 972, 934, 916, 776, 741, 684

* Measured in CDCl₃ with TMS as an internal standard unless otherwise specifically indicated.

Table 4

Comp'd No.	Structural formula	Property Melting point (recrystallization solvent)	NMR (δ ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
1 3		Colorless prism crystals 184.0-186.0°C (methanol)	(400MHz) (DMSO-d ₆ /TMS) 1.66 (2H, m), 1.84 (2H, m), 2.78 (2H, m), 2.96 (3H, s), 3.42 (2H, m), 3.63 (2H, t, J=6.6Hz), 3.96 (2H, t, J= 6.9Hz), 7.02 (1H, d, J=2.5Hz), 7.27 (1H, d, J=2.5Hz), 10.64 (1H, s)	(KBr) 3230, 1598, 1530, 1408, 1360, 1325, 1247, 1186, 1144, 1044, 968, 950, 917, 826, 794, 679
1 4		Colorless prism crystals 106.5-111.5°C (ethyl acetate- hexane)	(400MHz) 1.82 (2H, m), 1.98 (2H, m), 2.23 (2H, m), 3.09 (3H, s), 3.29 (1H, m), 3.54 (2H, dt, J=1.7Hz, 6.3Hz), 3.64 (1H, m), 3.97 (1H, m), 4.15 (1H, m), 4.91 (1H, m), 6.66 (1H, d, J=2.9Hz), 6.72 (1H, d, J= 2.9Hz)	(KBr) 3251, 2930, 2871, 1593, 1545, 1512, 1466, 1390, 1316, 1292, 1258, 1211, 1102, 1057, 962, 722
1 5		Brown oil ----- Hydrochloride Pale yellow prism crystals 225°C (decomp'd) (methanol- isopropyl ether)	(270MHz) 1.62 (2H, m), 1.79 (2H, m), 1.75-2.45 (6H, m), 2.50 (2H, m), 2.84 (2H, m), 3.00 (2H, m), 3.22 (3H, s), 3.29 (1H, m), 3.66 (2H, m), 4.33 (2H, m), 6.78 (1H, d, J= 2.6Hz), 6.91 (1H, d, J=2.6Hz), 7.14 (2H, t, J=8.6Hz), 7.95 (2H, d d, J= 5.3Hz, 8.6Hz),	(CHCl ₃) 2940, 2800, 1680, 1650, 1620, 1600, 1500, 1410, 1390, 1050, 975, 910
1 6		Pale yellow prism crystals 93.0-95.0°C (ethyl acetate- hexane)	(400MHz) 1.55 (2H, m), 1.80 (2H, m), 2.40 (2H, m), 2.58 (4H, m), 2.84 (2H, m), 3.19 (4H, m), 3.21 (3H, s), 3.65 (2H, m), 4.34 (2H, t, J= 7.2Hz), 6.78 (1H, d, J=2.6Hz), 6.85 (1H, m), 6.88-6.95 (3H, m), 7.26 (2H, m)	(KBr) 3089, 2811, 1634, 1499, 1407, 1384, 1309, 1227, 1138, 926, 804, 758, 695



* Measured in CDCl₃ with TMS as an internal standard unless otherwise specifically indicated.

Table 5

Comp'd No.	Structural formula	Property Melting point (recrystallization solvent)	NMR (δ ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
17		Pale yellow powdery crystals 114.0-117.0°C (2-propanol)	(400MHz) 1.55(2H, m), 1.80(2H, m), 2.00-2.17(6H, m), 2.40(2H, m), 2.85(2H, m), 2.98-3.12(3H, m), 3.22(3H, s), 3.65(2H, m), 4.34(2H, t, J=7.2Hz), 6.79(1H, d, J=2.6Hz), 6.91(1H, d, J=2.6Hz), 7.05(1H, d t, J=2.1Hz, 8.9Hz), 7.23(1H, d d, J=2.1Hz, 8.5Hz), 7.69(1H, d d, J=5.1Hz, 8.7Hz)	(KBr) 3098, 2944, 1652, 1616, 1502, 1395, 1316, 1269, 1250, 1137, 960, 905, 830, 758
18		Colorless powdery crystals 173.5-175.5°C (ethanol-ether)	(400MHz) 1.49(2H, m), 1.75-1.90(6H, m), 2.07(2H, m), 2.35(2H, m), 2.80(2H, m), 2.93(2H, m), 3.14-3.22(4H, m), 3.65(2H, m), 3.93(2H, t, J=7.1Hz), 7.13(2H, m), 7.33(2H, m), 7.95(2H, m)	(KBr) 3120, 2945, 1670, 1608, 1541, 1519, 1445, 1394, 1323, 1248, 1210, 1177, 1144, 972, 860, 762
19		Colorless oil	(400MHz) 1.54(2H, m), 1.77(2H, m), 1.82-1.90(4H, m), 2.14(2H, m), 2.39(2H, t, J=7.4Hz), 2.97(2H, m), 3.02(2H, m), 3.12(3H, s), 3.22(1H, quint, J=7.3Hz), 3.50(2H, m), 4.16(2H, t, J=7.2Hz), 6.65(1H, d, J=2.8Hz), 6.69(1H, d, J=2.8Hz), 7.13(2H, m), 7.95(2H, m)	(KBr) 2945, 1680, 1598, 1505, 1397, 1244, 1209, 1158, 974, 937, 854, 736
20		Colorless prism crystals 200.0-203.0°C (methanol-chloroform)	(270MHz) 1.55(2H, m), 1.74(2H, m), 2.43(2H, m), 2.64(4H, m), 2.98(2H, m), 3.10(3H, s), 3.22(4H, m), 3.48(2H, m), 4.14(2H, m), 6.64(1H, d, J=2.6Hz), 6.67(1H, d, J=2.6Hz), 6.82-6.96(3H, m), 7.26(2H, m), 10.95(1H, br. s)	(CHCl3) 3570, 2940, 2825, 1620, 1600, 1500, 1395, 1090, 970, 910

* Measured in CDCl₃ with TMS as an internal standard unless otherwise specifically indicated.

Table 6

Comp'd No.	Structural formula	Property Melting point (recrystallization solvent)	NMR (δ ppm)* (): observation frequency	IR (cm^{-1}) (): measuring method
2 1		Colorless powdery crystals 172.0-178.0°C (chloroform)	(400MHz) 1.51 (2H, m), 1.75-1.90 (6H, m), 2.07 (2H, m), 2.36 (2H, m), 2.90-3.00 (4H, m), 3.11 (3H, s), 3.18 (1H, m), 3.53 (2H, m), 3.89 (2H, t, J=7.1Hz), 6.95 (1H, d, J= 2.5Hz), 7.13 (2H, m), 7.27 (1H, d, J= 2.5Hz), 7.84 (1H, br. s), 7.95 (2H, m)	(KBr) 2942, 1677, 1598, 1537, 1508, 1410, 1324, 1215, 1142, 1039, 969, 913, 856, 796
2 2		Colorless prism crystals 173.0°C-175.5°C (chloroform-ether)	(400MHz) 1.59 (2H, m), 1.74-2.07 (8H, m), 2.12- 2.36 (4H, m), 2.82 (1H, m), 2.95 (1H, m), 3.13 (3H, s), 3.18 (1H, m), 3.30 (1H, m), 3.75 (1H, m), 4.01 (2H, t, J=7.4Hz), 4.92 (1H, t, J=4.4Hz), 6.63 (1H, d, J=2.9Hz), 6.74 (1H, d, J=2.9Hz), 7.13 (2H, m), 7.94 (2H, m),	(KBr) 3274, 2935, 2808, 1672, 1586, 1511, 1317, 1284, 1207, 1154, 1101, 1057, 973, 856, 740

* Measured in CDCl_3 with TMS as an internal standard unless otherwise specifically indicated.

- 66 -

Tests

With respect to the compounds of the present invention, their anti- α_1 action and anti-serotonin (5-HT) action were investigated by the methods which will be described below. The results of some representative compounds are shown in Table 7.

(1) Anti- α_1 action

The thoracic aorta of each Hartley male guinea pig (body weight: 300-500 g) was excised. A preparation cut in a helical form was suspended under 1 g load in a Magnus cylinder filled with the Tyrode solution which had been aerated with a gas mixture of 95% O₂ and 5% CO₂ and maintained at 37°C. Using an isometric transducer ("TB-612J", manufactured by Nihon Kohden Corporation) and a pressure preamplifier ("AP-620G", manufactured by Nihon Kohden Corporation), variations in tension were measured. The isometric tensions were recorded on a thermal pen-writing recorder ("WT-647G", manufactured by Nihon Kohden Corporation). Taking the tonic contraction induced by 10⁻⁵ M norepinephrine (NE) as 100%, the percent contractions upon addition of each test drug at 10⁻⁸ M and 10⁻⁷ M were determined as anti- α_1 action.

(2) Anti-serotonin (5-HT) action

The superior mesenteric artery of each Hartley

- 67 -

male guinea pig (body weight: 300-500 g) was excised. A preparation cut in a helical form was suspended under resting tension of 0.3 g in a Magnus cylinder filled with the Tyrode solution which had been aerated with a gas mixture of 95% O₂ and 5% CO₂ and maintained at 37°C. Using an isometric transducer ("UL-10", manufactured by SHINKOH K.K.) and a pressure preamplifier ("DSA-605A", manufactured by SHINKOH K.K.), variations in tension were measured. The isometric tensions were recorded on a pen-writing recorder ("VP-6537A", manufactured by NATIONAL K.K.). Taking the contraction induced by 10⁻⁵ M serotonin (5-HT) as 100%, the percent contractions by 10⁻⁵ M 5-HT in the presence of each test drug at 10⁻⁸ M, 10⁻⁷ M and 10⁻⁶ M were determined as anti-5-HT action.

(3) Results

Table 7

Comp'd No.	Anti α_1 action (% of Control)		Anti 5-HT action (% of Control)		
	10 ⁻⁸ M	10 ⁻⁷ M	10 ⁻⁸ M	10 ⁻⁷ M	10 ⁻⁶ M
15	59.9	35.4	NT	75.3	26.6
17	27.8	14.0	80.7	55.2	8.9
18	76.0	37.3	NT	81.7	25.3
21	69.7	36.9	NT	85.5	24.9

NT ... Not tested.

- 68 -


Capability of Exploitation in Industry

The pyrroloazepine derivatives (I) and their salts according to the present invention have strong α_1 -blocking action and serotonin-2 blocking action, and also have high safety. Accordingly, the present invention has made it possible to provide preventives or therapeutics for general circulatory diseases such as hypertension, heart failure, ischemic heart diseases, cerebrovascular disturbances and peripheral circulatory disturbances.

- 69 -

CLAIMS

1 1. A pyrroloazepine compound or a salt thereof,
2 said pyrroloazepine compound being represented by the
3 following formula (I):




```

5      wherein
6          the ring P represented by

```



8 means a pyrrole ring represented by the following
9 structure:




11 in which A represents an alkylene group, an alkenylene
12 group or an alkynylene group, and Y represents a group



14 in which W represents CH, C= or a nitrogen atom; and,

- 70 -

when W represents CH, m stands for 0 or 1, B represents an oxygen atom, a sulfur atom, a carbonyl group, a sulfinyl group, a sulfonyl group, an alkylene group, an alkenylene group, a group $-C(OH)R_1-$ in which R_1 represents a substituted or unsubstituted aryl group, a group $-CHR_2-$ in which R_2 represents a substituted or unsubstituted aryl group, or a substituted or unsubstituted, cyclic or acyclic acetal group; when W represents C=, m stands for 1, B represents a group



in which the double bond is coupled with W and R_3 represents a substituted or unsubstituted aryl group or a substituted or unsubstituted aralkyl group; when W represents a nitrogen atom, m stands for 0 or 1, and B represents a carbonyl group, a sulfonyl group, an alkylene group, an alkenylene group or a group $-CHR_4-$ in which R_4 represents a substituted or unsubstituted aryl group; E_1 and E_2 each independently represents a hydrogen atom or a lower alkyl group; and D represents a substituted or unsubstituted aromatic hydrocarbon group or a substituted or unsubstituted aromatic heterocyclic group;

the dashed line indicates the presence or absence of a bond; and, when the bond indicated by the dashed

- 71 -

39 line is present, Z_2 is not present and Z_1 represents a
 40 hydrogen atom but, when the bond indicated by the
 41 dashed line is absent, Z_1 and Z_2 both represent
 42 hydrogen atoms; Z_1 represents a hydrogen atom and Z_2
 43 represents a hydroxyl group; Z_1 and Z_2 both represent
 44 groups SR_5 in which R_5 represents a substituted or un-
 45 substituted alkyl group, a substituted or unsubstituted
 46 aralkyl group or a substituted or unsubstituted aryl
 47 group; or Z_1 and Z_2 are combined together to represent
 48 an oxygen atom, a group NOR_6 in which R_6 represents a
 49 hydrogen atom, a substituted or unsubstituted alkyl
 50 group, a substituted or unsubstituted aralkyl group or
 51 a substituted or unsubstituted aryl group, or a group

52



53 in which G represents a substituted or unsubstituted
 54 ethylene group or a substituted or unsubstituted
 55 trimethylene group; and

56 R represents a hydrogen atom, a linear or
 57 branched alkyl group, a cycloalkyl group, a cycloalkyl-
 58 alkyl group or a substituted or unsubstituted aralkyl
 59 group.

1 2. A pyrroloazepine compound or a salt thereof
 2 according to claim 1, wherein in the formula (I), Z_1

- 72 -

3 and Z_2 are combined together to represent an oxygen
4 atom or the group NOR_6 .

1 3. A pyrroloazepine compound or a salt thereof
2 according to claim 1, wherein in the formula (I), Z_1
3 represents a hydrogen atom and Z_2 represents a hydroxyl
4 group.

1 4. A pyrroloazepine compound or a salt thereof
2 according to claim 1, 2 or 3, wherein in the formula
3 (I), A represents a tetramethylene group.

1 5. A pyrroloazepine compound or a salt thereof
2 according to any one of claims 1-4, wherein in the for-
3 mula (I), W represents CH, B represents a carbonyl
4 group, m stands for 1, and D represents a substituted
5 or unsubstituted phenyl group.


1 6. A pyrroloazepine compound or a salt thereof
2 according to any one of claims 1-4, wherein in the for-
3 mula (I), W represents CH, m stands for 0, and D
4 represents a substituted or unsubstituted aromatic
5 hydrocarbon group or a substituted or unsubstituted
6 aromatic heterocyclic group.

1 7. A pyrroloazepine compound or a salt thereof
2 according to any one of claims 1-6, wherein in the for-
3 mula (I), E_1 and E_2 both represent hydrogen atoms.

1 8. A process for the preparation of a pyrrolo-
2 azepine compound represented by the following formula


- 73 -

3 (Ia) or (Ia'):




5 wherein A, R and Y have the same meanings as defined
6 above, which comprises:

7 reacting a compound, which is represented by the
8 following formula (III):




10 wherein A has the same meaning as defined above and X
11 and X' represent the same or different eliminative
12 groups, to a compound represented by the following for-
13 mula (II) or (II'):



15 wherein R has the same meaning as defined above, there-
16 by obtaining a compound represented by the following
17 formula (IV) or (IV'):


- 74 -

18



(IV)


or



(IV')

19 wherein A, R and X have the same meanings as defined
 20 above; and then


21 reacting a nitrogen-containing compound, which is
 22 represented by the following formula (V):



24 wherein Y has the same meaning as defined above, to the
 25 compound represented by the formula (IV) or (IV').


1 9. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ia) or (Ia'):

4



(Ia)

or



(Ia')

5 wherein A, R and Y have the same meanings as defined
 6 above, which comprises:


7 reacting a compound, which is represented by the

- 75 -

8 following formula (VI):


9 $X-A-Y$ (VI)

10 wherein A, X and Y have the same meanings as defined
11 above, to a compound represented by the following for-
12 mula (II) or (II'):



14 wherein R has the same meaning as defined above.


1 10. A process for the preparation of a pyrrolo-
2 azepine compound represented by the following formula
3 (Ic) or (Ic'):



5 wherein Z₁' and Z₂' both represent groups -SR₅ in which
6 R₅ has the same meaning as defined above or are com-
7 bined together to represent a group -S-G-S in which G
8 has the same meaning as defined above, Y' represents
9 a group


- 76 -

10



11 in which, when W represents CH, B' represents an oxygen
 12 atom, a sulfur atom, a sulfinyl group, a sulfonyl
 13 group, an alkylene group, an alkenylene group, a group
 14 -C(OH)R₁- in which R₁ represents a substituted or un-
 15 substituted aryl group, a group -CHR₂- in which R₂
 16 represents a substituted or unsubstituted aryl group,
 17 or a substituted or unsubstituted, cyclic or acyclic
 18 acetal group; when W represents C=, B' represents a
 19 group


20




21 in which the double bond is coupled with W and R₃
 22 represents a substituted or unsubstituted aryl group or
 23 a substituted or unsubstituted aralkyl group; when W
 24 represents a nitrogen atom, B' represents a carbonyl
 25 group, a sulfonyl group, an alkylene group, an
 26 alkenylene group or a group -CHR₄- in which R₄
 27 represents a substituted or unsubstituted aryl group;
 28 and D, E₁, E₂ and m have the same meanings as defined
 29 above, and A and R have the same meanings as defined
 30 above, which comprises:

- 77 -

reacting a pyrroloazepine compound, which is represented by the following formula (Ib) or (Ib'):




wherein Y', A and R have the same meanings as defined above, with a compound represented by the following formula (VIIa) or (VIIb):



wherein G and R₅ have the same meanings as defined above.

11. A process for the preparation of a pyrroloazepine compound represented by the following formula (Id) or (Id'):



wherein A, R, R₆ and Y' have the same meanings as


- 78 -

defined above, which comprises:

reacting a hydroxylamine or a derivative thereof
represented by the following formula (VIII):




wherein R_6 has the same meaning as defined above, to a
pyrroloazepine compound represented by the following
formula (Ib) or (Ib'):



(Ib)


or



(Ib')


wherein A, R and Y' have the same meanings as defined
above.

12. A process for the preparation of a pyrrolo-
azepine compound represented by the following formula
(Ie) or (Ie'):



(Ie)

or




(Ie')

wherein A, R and Y' have the same meanings as defined

- 79 -


6 above, which comprises:

7 reducing a pyrroloazepine compound represented by
8 the following formula (Ib) or (Ib'):



10 wherein A, R and Y' have the same meanings as defined
11 above.

1 13. A process for the preparation of a pyrrolo-
2 azepine compound represented by the following formula
3 (If) or (If'):




5 wherein A, R, Y, Z₁' and Z₂' have the same meanings as
6 defined above, which comprises:


7 reacting a compound represented by the following
8 formula (VIIa) or (VIIb):

- 80 -


9



10 wherein G and R₅ have the same meanings as defined
11 above, to a compound represented by the following for-
12 mula (II) or (II'):



14 wherein R has the same meaning as defined above, there-
15 by obtaining a compound represented by the following
16 formula (IX) or (IX'):




18 wherein R, Z₁' and Z₂' have the same meanings as
19 defined above; and then
20 reacting a nitrogen-containing compound, which is
21 represented by the following formula (VI):
22
$$X-A-Y \quad (VI)$$

23 wherein A, X and Y have the same meanings as defined
24 above, to the compound represented by the formula (IX)

- 81 -


25 or (IX').

1 14. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ig) or (Ig'):




5 wherein A, R, R₆ and Y have the same meanings as
 6 defined above, which comprises:

7 reacting a hydroxylamine or a derivative thereof,
 8 which is represented by the following formula (VIII):




10 wherein R₆ has the same meaning as defined above, to a
 11 compound represented by the following formula (IV) or
 12 (IV'):



14 wherein A, R and X have the same meanings as defined

- 82 -

15 above, thereby obtaining a compound represented by the
 16 following formula (X) or (X'):




18 wherein A, R, R₆ and X have the same meanings as
 19 defined above; and then

20 reacting a nitrogen-containing compound, which is
 21 represented by the following formula (V):



23 wherein Y has the same meanings as defined above, to
 24 the compound represented by the formula (X) or (X').

1 15. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ih) or (Ih'):




5 wherein A, R and Y have the same meanings as defined

- 83 -


above, which comprises:

reducing a compound represented by the following
formula (IV) or (IV'):




(IV)

or




(IV')

wherein A, R and X have the same meanings as defined
above, thereby obtaining a compound represented by the
following formula (XI) or (XI'):



(XI)

or



(XI')

wherein A, R and X have the same meanings as defined
above; and then

reacting a nitrogen-containing compound
represented by the following formula (V):




(V)


wherein Y has the same meaning as defined above, to the
compound represented by the formula (XI) or (XI').

- 84 -

1 16. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ii) or (Ii'):




(Ii)




(Ii')

5 wherein A, R and Y have the same meanings as defined
 6 above, which comprises:

7 subjecting a compound, which is represented by
 8 the following formula (XI) or (XI'):



(XI)




(XI')

10 wherein A, R and X have the same meanings as defined
 11 above, to a dehydrating reaction, thereby obtaining a
 12 compound represented by the following formula (XII) or
 13 (XII'):


- 85 -

14



(XII)

or



(XII')

15 wherein A, R and X have the same meanings as defined
 16 above; and then


17 reacting a nitrogen-containing compound
 18 represented by the following formula (V):

19
$$\text{H-Y} \quad \quad \quad \text{(V)}$$

20 wherein Y has the same meaning as defined above, to the
 21 compound represented by the formula (XII) or (XII').


1 17. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ii) or (Ii'):

4



(Ii)

or




(Ii')

5 wherein A, R and Y have the same meanings as defined
 6 above, which comprises:

7 subjecting a compound, which is represented by


- 86 -

8 the following formula (Ih) or (Ih'):



10 wherein A, R and Y have the same meanings as defined
 11 above, to a dehydrating reaction.


1 18. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (Ij) or (Ij'):




5 wherein A, R and Y have the same meanings as defined
 6 above, which comprises:

7 reducing a compound represented by the following
 8 formula (XII) or (XII'):

- 87 -




10 wherein A, R and X have the same meanings as defined
11 above, thereby obtaining a compound represented by the
12 following formula (XIII) or (XIII'):



14 wherein A, R and X have the same meanings as defined
15 above; and then


16 reacting a nitrogen-containing compound
17 represented by the following formula (V):



19 wherein Y has the same meaning as defined above, to the
20 compound represented by the formula (XIII) or (XIII').


1 19. An intermediate for the production of a
2 pharmaceutical, which is represented by the following
3 formula (IIa) or (IIa'):

- 88 -



5 wherein R' represents a linear or branched alkyl group,
 6 a cycloalkyl group, a cycloalkyl-alkyl group or a sub-
 7 stituted or unsubstituted aralkyl group.


1 20. An intermediate for the production of a
 2 pharmaceutical, which is represented by the following
 3 formula (XIV) or (XIV'):



5 wherein the dashed line, A, R, X, Z₁ and Z₂ have the
 6 same meanings as defined above.


1 21. A process for the preparation of a pyrrolo-
 2 azepine compound represented by the following formula
 3 (IIa) or (IIa'):

- 89 -



5 wherein R' has the same meaning as defined above, which
6 comprises:

7 reacting a β -aminopropionic acid or a derivative
8 thereof, which is represented by the following formula
9 (XVI):




11 wherein R₇ represents a hydrogen atom or a carboxyl-
12 protecting group and R' has the same meaning as defined
13 above, to a pyrrole-3-carboxylic acid or a derivative
14 thereof represented by the following formula (XV):



16 wherein Q represents a hydroxyl group, an alkoxy group
17 or an eliminative group easily replaceable by an amino
18 group, thereby obtaining a compound represented by the
19 following formula (XVII):

- 90 -

20




21 wherein R' and R₇ have the same meanings as defined
 22 above; and then

23 subjecting the compound, which is represented by
 24 the formula (XVII), to ring closure.

1 22. An intermediate for the production of a
 2 pharmaceutical, which is represented by the following
 3 formula (XVII):

4



5 wherein R' and R₇ have the same meanings as defined
 6 above.

1 23. A pharmaceutical comprising, as an effective
 2 ingredient, a pyrroloazepine compound or a salt thereof
 3 according to claim 1.

1 24. A therapeutic for circulatory diseases, com-
 2 prising as an effective ingredient a pyrroloazepine
 3 compound or a salt thereof according to claim 1.

1 25. A serotonin-2 receptor antagonist, comprising
 2 as an effective ingredient a pyrroloazepine compound or

- 91 -

3 a salt thereof according to claim 1.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 98/01085

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D487/04 C07D207/40 A61K31/55 //(C07D487/04,223:00,
209:00)

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 441 349 A (SUNTORY) 14 August 1991 see page 48 ---	1,23
A	EP 0 551 527 A (SUNTORY) 21 July 1993 see page 38, line 1 - line 39; claim 1 ---	1,23
P, A	EP 0 807 632 A (SUNTORY) 19 November 1997 see page 82, line 50 - page 83, line 45; claim 1 -----	1,23

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 June 1998

Date of mailing of the international search report

24/06/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 98/01085

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 441349 A	14-08-1991	JP 5097856 A	20-04-1993
		AT 132497 T	15-01-1996
		AU 642960 B	04-11-1993
		AU 7080691 A	08-08-1991
		CA 2035749 A	08-08-1991
		DE 69115943 D	15-02-1996
		DK 441349 T	13-05-1996
		ES 2084719 T	16-05-1996
		US 5416082 A	16-05-1995
		US 5206239 A	27-04-1993
		US 5391731 A	21-02-1995
EP 551527 A	21-07-1993	AU 655841 B	12-01-1995
		AU 2403192 A	02-03-1993
		CA 2093631 A	08-02-1993
		WO 9303031 A	18-02-1993
		US 5397780 A	14-03-1995
EP 807632 A	19-11-1997	AU 7655896 A	27-06-1997
		CA 2212092 A	12-06-1997
		WO 9720845 A	12-06-1997